

Chapter udf

Natural Deduction

This chapter presents a natural deduction system in the style of Gentzen/Prawitz.

To include or exclude material relevant to natural deduction as a proof system, use the “prfND” tag.

ntd.1 Rules and Derivations

fol:ntd:rul:
sec

Natural deduction systems are meant to closely parallel the informal reasoning used in mathematical proof (hence it is somewhat “natural”). Natural deduction proofs begin with assumptions. Inference rules are then applied. Assumptions are “discharged” by the \neg Intro, \rightarrow Intro, \forall Elim and \exists Elim inference rules, and the label of the discharged assumption is placed beside the inference for clarity.

explanation

Definition ntd.1 (Assumption). An *assumption* is any sentence in the top-most position of any branch.

Derivations in natural deduction are certain trees of sentences, where the topmost sentences are assumptions, and if a sentence stands below one, two, or three other sequents, it must follow correctly by a rule of inference. The sentences at the top of the inference are called the *premises* and the sentence below the *conclusion* of the inference. The rules come in pairs, an introduction and an elimination rule for each logical operator. They introduce a logical operator in the conclusion or remove a logical operator from a premise of the rule. Some of the rules allow an assumption of a certain type to be discharged. To indicate which assumption is discharged by which inference, we also assign labels to both the assumption and the inference. This is indicated by writing the assumption as “[φ]ⁿ.”

It is customary to consider rules for all the logical operators \wedge , \vee , \rightarrow , \neg , and \perp , even if some of those are defined.

ntd.2 Propositional Rules

Rules for \wedge

fol:ntd:prl:
sec

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge\text{Intro} \qquad \frac{\varphi \wedge \psi}{\varphi} \wedge\text{Elim}$$

$$\frac{\varphi \wedge \psi}{\psi} \wedge\text{Elim}$$

Rules for \vee

$$\frac{\varphi}{\varphi \vee \psi} \vee\text{Intro}$$

$$\frac{\psi}{\varphi \vee \psi} \vee\text{Intro}$$

$$n \frac{\varphi \vee \psi}{\chi} \vee\text{Elim}$$

$$\begin{array}{c} [\varphi]^n \\ \vdots \\ \chi \\ \hline \chi \end{array} \quad \begin{array}{c} [\psi]^n \\ \vdots \\ \chi \\ \hline \chi \end{array} \vee\text{Elim}$$

Rules for \rightarrow

$$n \frac{\begin{array}{c} [\varphi]^n \\ \vdots \\ \psi \end{array}}{\varphi \rightarrow \psi} \rightarrow\text{Intro}$$

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow\text{Elim}$$

Rules for \neg

$$n \frac{\begin{array}{c} [\varphi]^n \\ \vdots \\ \perp \end{array}}{\neg\varphi} \neg\text{Intro}$$

$$\frac{\neg\varphi \quad \varphi}{\perp} \neg\text{Elim}$$

Rules for \perp

$$\frac{\perp}{\varphi} \perp_I \qquad \begin{array}{c} [\neg\varphi]^n \\ \vdots \\ \vdots \\ n \frac{\perp}{\varphi} \perp_C \end{array}$$

Note that \neg -Intro and \perp_C are very similar: The difference is that \neg -Intro derives a negated **sentence** $\neg\varphi$ but \perp_C a positive **sentence** φ .

Whenever a rule indicates that some assumption may be discharged, we take this to be a permission, but not a requirement. E.g., in the \rightarrow -Intro rule, we may discharge any number of assumptions of the form φ in the **derivation** of the premise ψ , including zero.

ntd.3 Quantifier Rules

fol:ntd:qrl:
sec

Rules for \forall

$$\frac{\varphi(a)}{\forall x \varphi(x)} \forall\text{Intro} \qquad \frac{\forall x \varphi(x)}{\varphi(t)} \forall\text{Elim}$$

In the rules for \forall , t is a closed term (a term that does not contain any variables), and a is a **constant symbol** which does not occur in the conclusion $\forall x \varphi(x)$, or in any assumption which is **undischarged** in the **derivation** ending with the premise $\varphi(a)$. We call a the *eigenvariable* of the \forall -Intro inference.¹

Rules for \exists

$$\frac{\varphi(t)}{\exists x \varphi(x)} \exists\text{Intro} \qquad \begin{array}{c} [\varphi(a)]^n \\ \vdots \\ \vdots \\ n \frac{\exists x \varphi(x)}{\chi} \exists\text{Elim} \end{array}$$

Again, t is a closed term, and a is a **constant symbol** which does not occur in the premise $\exists x \varphi(x)$, in the conclusion χ , or any assumption which is **undischarged**

¹We use the term “eigenvariable” even though a in the above rule is a constant. This has historical reasons.

in the **derivations** ending with the two premises (other than the assumptions $\varphi(a)$). We call a the *eigenvariable* of the \exists Elim inference.

The condition that an eigenvariable neither occur in the premises nor in any assumption that is **undischarged** in the **derivations** leading to the premises for the \forall Intro or \exists Elim inference is called the *eigenvariable condition*.

explanation Recall the convention that when φ is a **formula** with the **variable** x free, we indicate this by writing $\varphi(x)$. In the same context, $\varphi(t)$ then is short for $\varphi[t/x]$. So we could also write the \exists Intro rule as:

$$\frac{\varphi[t/x]}{\exists x \varphi} \exists\text{Intro}$$

Note that t may already occur in φ , e.g., φ might be $P(t, x)$. Thus, inferring $\exists x P(t, x)$ from $P(t, t)$ is a correct application of \exists Intro—you may “replace” one or more, and not necessarily all, occurrences of t in the premise by the bound **variable** x . However, the eigenvariable conditions in \forall Intro and \exists Elim require that the **constant symbol** a does not occur in φ . So, you cannot correctly infer $\forall x P(a, x)$ from $P(a, a)$ using \forall Intro.

explanation In \exists Intro and \forall Elim there are no restrictions, and the term t can be anything, so we do not have to worry about any conditions. On the other hand, in the \exists Elim and \forall Intro rules, the eigenvariable condition requires that the **constant symbol** a does not occur anywhere in the conclusion or in an **undischarged** assumption. The condition is necessary to ensure that the system is sound, i.e., only **derives sentences** from **undischarged** assumptions from which they follow. Without this condition, the following would be allowed:

$$\frac{\exists x \varphi(x) \quad \frac{[\varphi(a)]^1}{\forall x \varphi(x)} * \forall\text{Intro}}{\forall x \varphi(x)} \exists\text{Elim}$$

However, $\exists x \varphi(x) \not\equiv \forall x \varphi(x)$.

As the elimination rules for quantifiers only allow substituting closed terms for **variables**, it follows that any **formula** that can be derived from a set of **sentences** is itself a **sentence**.

ntd.4 Derivations

explanation We’ve said what an assumption is, and we’ve given the rules of inference. **Derivations** in natural deduction are inductively generated from these: each **derivation** either is an assumption on its own, or consists of one, two, or three **derivations** followed by a correct inference. **fol:ntd:der:sec**

Definition ntd.2 (Derivation). A *derivation* of a **sentence** φ from assumptions Γ is a finite tree of **sentences** satisfying the following conditions:

1. The topmost **sentences** of the tree are either in Γ or are **discharged** by an inference in the tree.

2. The bottommost **sentence** of the tree is φ .
3. Every **sentence** in the tree except the sentence φ at the bottom is a premise of a correct application of an inference rule whose conclusion stands directly below that **sentence** in the tree.

We then say that φ is the *conclusion* of the **derivation** and Γ its **undischarged** assumptions.

If a **derivation** of φ from Γ exists, we say that φ is *derivable* from Γ , or in symbols: $\Gamma \vdash \varphi$. If there is a **derivation** of φ in which every assumption is **discharged**, we write $\vdash \varphi$.

Example ntd.3. Every assumption on its own is a **derivation**. So, e.g., φ by itself is a **derivation**, and so is ψ by itself. We can obtain a new **derivation** from these by applying, say, the \wedge Intro rule,

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge\text{Intro}$$

These rules are meant to be general: we can replace the φ and ψ in it with any **sentences**, e.g., by χ and θ . Then the conclusion would be $\chi \wedge \theta$, and so

$$\frac{\chi \quad \theta}{\chi \wedge \theta} \wedge\text{Intro}$$

is a correct **derivation**. Of course, we can also switch the assumptions, so that θ plays the role of φ and χ that of ψ . Thus,

$$\frac{\theta \quad \chi}{\theta \wedge \chi} \wedge\text{Intro}$$

is also a correct **derivation**.

We can now apply another rule, say, \rightarrow Intro, which allows us to conclude a conditional and allows us to **discharge** any assumption that is identical to the antecedent of that conditional. So both of the following would be correct **derivations**:

$$1 \frac{\frac{[\chi]^1 \quad \theta}{\chi \wedge \theta} \wedge\text{Intro}}{\chi \rightarrow (\chi \wedge \theta)} \rightarrow\text{Intro} \quad 1 \frac{\frac{\chi \quad [\theta]^1}{\chi \wedge \theta} \wedge\text{Intro}}{\theta \rightarrow (\chi \wedge \theta)} \rightarrow\text{Intro}$$

They show, respectively, that $\theta \vdash \chi \rightarrow (\chi \wedge \theta)$ and $\chi \vdash \theta \rightarrow (\chi \wedge \theta)$.

Remember that discharging of assumptions is a permission, not a requirement: we don't have to discharge the assumptions. In particular, we can apply a rule even if the assumptions are not present in the **derivation**. For instance, the following is legal, even though there is no assumption φ to be **discharged**:

$$1 \frac{\psi}{\varphi \rightarrow \psi} \rightarrow\text{Intro}$$

ntd.5 Examples of Derivations

fol:ntd:pro:
sec

Example ntd.4. Let's give a **derivation** of the **sentence** $(\varphi \wedge \psi) \rightarrow \varphi$.

We begin by writing the desired conclusion at the bottom of the **derivation**.

$$\frac{}{(\varphi \wedge \psi) \rightarrow \varphi}$$

Next, we need to figure out what kind of inference could result in a **sentence** of this form. The **main operator** of the conclusion is \rightarrow , so we'll try to arrive at the conclusion using the \rightarrow Intro rule. It is best to write down the assumptions involved and label the inference rules as you progress, so it is easy to see whether all assumptions have been **discharged** at the end of the proof.

$$1 \frac{\begin{array}{c} [\varphi \wedge \psi]^1 \\ \vdots \\ \vdots \\ \varphi \end{array}}{(\varphi \wedge \psi) \rightarrow \varphi} \rightarrow\text{Intro}$$

We now need to fill in the steps from the assumption $\varphi \wedge \psi$ to φ . Since we only have one connective to deal with, \wedge , we must use the \wedge elim rule. This gives us the following proof:

$$1 \frac{\frac{[\varphi \wedge \psi]^1}{\varphi} \wedge\text{Elim}}{(\varphi \wedge \psi) \rightarrow \varphi} \rightarrow\text{Intro}$$

We now have a correct **derivation** of $(\varphi \wedge \psi) \rightarrow \varphi$.

Example ntd.5. Now let's give a **derivation** of $(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)$.

We begin by writing the desired conclusion at the bottom of the **derivation**.

$$\frac{}{(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)}$$

To find a logical rule that could give us this conclusion, we look at the logical connectives in the conclusion: \neg , \vee , and \rightarrow . We only care at the moment about the first occurrence of \rightarrow because it is the **main operator** of the **sentence** in the end-sequent, while \neg , \vee and the second occurrence of \rightarrow are inside the scope of another connective, so we will take care of those later. We therefore start with the \rightarrow Intro rule. A correct application must look like this:

$$1 \frac{\begin{array}{c} [\neg\varphi \vee \psi]^1 \\ \vdots \\ \vdots \\ \varphi \rightarrow \psi \end{array}}{(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)} \rightarrow\text{Intro}$$

This leaves us with two possibilities to continue. Either we can keep working from the bottom up and look for another application of the \rightarrow Intro rule, or we can work from the top down and apply a \vee Elim rule. Let us apply the latter. We will use the assumption $\neg\varphi \vee \psi$ as the leftmost premise of \vee Elim. For a valid application of \vee Elim, the other two premises must be identical to the conclusion $\varphi \rightarrow \psi$, but each may be derived in turn from another assumption, namely one of the two disjuncts of $\neg\varphi \vee \psi$. So our **derivation** will look like this:

$$\begin{array}{c}
 \begin{array}{c} [\neg\varphi]^2 \\ \vdots \\ \varphi \rightarrow \psi \end{array} \quad \begin{array}{c} [\psi]^2 \\ \vdots \\ \varphi \rightarrow \psi \end{array} \\
 \hline
 2 \frac{[\neg\varphi \vee \psi]^1 \quad \varphi \rightarrow \psi \quad \varphi \rightarrow \psi}{\varphi \rightarrow \psi} \vee\text{Elim} \\
 \hline
 1 \frac{\varphi \rightarrow \psi}{(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)} \rightarrow\text{Intro}
 \end{array}$$

In each of the two branches on the right, we want to **derive** $\varphi \rightarrow \psi$, which is best done using \rightarrow Intro.

$$\begin{array}{c}
 \begin{array}{c} [\neg\varphi]^2, [\varphi]^3 \\ \vdots \\ \psi \end{array} \quad \begin{array}{c} [\psi]^2, [\varphi]^4 \\ \vdots \\ \psi \end{array} \\
 \hline
 2 \frac{[\neg\varphi \vee \psi]^1 \quad 3 \frac{\psi}{\varphi \rightarrow \psi} \rightarrow\text{Intro} \quad 4 \frac{\psi}{\varphi \rightarrow \psi} \rightarrow\text{Intro}}{\varphi \rightarrow \psi} \vee\text{Elim} \\
 \hline
 1 \frac{\varphi \rightarrow \psi}{(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)} \rightarrow\text{Intro}
 \end{array}$$

For the two missing parts of the **derivation**, we need **derivations** of ψ from $\neg\varphi$ and φ in the middle, and from φ and ψ on the left. Let's take the former first. $\neg\varphi$ and φ are the two premises of \neg Elim:

$$\begin{array}{c}
 \frac{[\neg\varphi]^2 \quad [\varphi]^3}{\perp} \neg\text{Elim} \\
 \vdots \\
 \psi
 \end{array}$$

By using \perp_I , we can obtain ψ as a conclusion and complete the branch.

$$\begin{array}{c}
 \begin{array}{c} [\neg\varphi]^2 \quad [\varphi]^3 \\ \vdots \\ \perp \end{array} \quad \begin{array}{c} [\psi]^2, [\varphi]^4 \\ \vdots \\ \psi \end{array} \\
 \hline
 2 \frac{[\neg\varphi \vee \psi]^1 \quad 3 \frac{\frac{\perp}{\psi} \perp_I}{\varphi \rightarrow \psi} \rightarrow\text{Intro} \quad 4 \frac{\psi}{\varphi \rightarrow \psi} \rightarrow\text{Intro}}{\varphi \rightarrow \psi} \vee\text{Elim} \\
 \hline
 1 \frac{\varphi \rightarrow \psi}{(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)} \rightarrow\text{Intro}
 \end{array}$$

Let's now look at the rightmost branch. Here it's important to realize that the definition of **derivation** allows assumptions to be discharged but does not require them to be. In other words, if we can derive ψ from one of the assumptions φ and ψ without using the other, that's ok. And to **derive** ψ from ψ is trivial: ψ by itself is such a **derivation**, and no inferences are needed. So we can simply delete the assumption φ .

$$\begin{array}{c}
 \frac{[\neg\varphi]^2 \quad [\varphi]^3}{\perp} \neg\text{Elim} \\
 \frac{\perp}{\psi} \perp_I \\
 \frac{[\neg\varphi \vee \psi]^1 \quad \frac{\varphi \rightarrow \psi}{\psi} \rightarrow\text{Intro}}{\varphi \rightarrow \psi} \rightarrow\text{Intro} \\
 \frac{[\psi]^2}{\varphi \rightarrow \psi} \rightarrow\text{Intro} \\
 \frac{[\neg\varphi \vee \psi]^1 \quad \frac{\varphi \rightarrow \psi}{\varphi \rightarrow \psi} \rightarrow\text{Intro}}{(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)} \rightarrow\text{Intro} \\
 \frac{(\neg\varphi \vee \psi) \rightarrow (\varphi \rightarrow \psi)}{\varphi \rightarrow \psi} \vee\text{Elim}
 \end{array}$$

Note that in the finished **derivation**, the rightmost \rightarrow Intro inference does not actually discharge any assumptions.

Example ntd.6. So far we have not needed the \perp_C rule. It is special in that it allows us to discharge an assumption that isn't a sub-formula of the conclusion of the rule. It is closely related to the \perp_I rule. In fact, the \perp_I rule is a special case of the \perp_C rule—there is a logic called “intuitionistic logic” in which only \perp_I is allowed. The \perp_C rule is a last resort when nothing else works. For instance, suppose we want to **derive** $\varphi \vee \neg\varphi$. Our usual strategy would be to attempt to **derive** $\varphi \vee \neg\varphi$ using \vee Intro. But this would require us to **derive** either φ or $\neg\varphi$ from no assumptions, and this can't be done. \perp_C to the rescue!

$$\begin{array}{c}
 [\neg(\varphi \vee \neg\varphi)]^1 \\
 \vdots \\
 \vdots \\
 \vdots \\
 \perp \\
 \frac{\perp}{\varphi \vee \neg\varphi} \perp_C
 \end{array}$$

Now we're looking for a **derivation** of \perp from $\neg(\varphi \vee \neg\varphi)$. Since \perp is the conclusion of \neg Elim we might try that:

$$\begin{array}{c}
 [\neg(\varphi \vee \neg\varphi)]^1 \quad [\neg(\varphi \vee \neg\varphi)]^1 \\
 \vdots \quad \vdots \\
 \vdots \quad \vdots \\
 \vdots \quad \vdots \\
 \neg\varphi \quad \varphi \\
 \frac{\neg\varphi \quad \varphi}{\perp} \neg\text{Elim} \\
 \frac{\perp}{\varphi \vee \neg\varphi} \perp_C
 \end{array}$$

Our strategy for finding a **derivation** of $\neg\varphi$ calls for an application of \neg Intro:

$$\begin{array}{c}
[\neg(\varphi \vee \neg\varphi)]^1, [\varphi]^2 \\
\vdots \\
2 \frac{\perp}{\neg\varphi} \neg\text{Intro} \quad \quad \quad [\neg(\varphi \vee \neg\varphi)]^1 \\
\vdots \\
\varphi \neg\text{Elim} \\
\hline
1 \frac{\perp}{\varphi \vee \neg\varphi} \perp_C
\end{array}$$

Here, we can get \perp easily by applying $\neg\text{Elim}$ to the assumption $\neg(\varphi \vee \neg\varphi)$ and $\varphi \vee \neg\varphi$ which follows from our new assumption φ by $\vee\text{Intro}$:

$$\begin{array}{c}
[\neg(\varphi \vee \neg\varphi)]^1 \quad \frac{[\varphi]^2}{\varphi \vee \neg\varphi} \vee\text{Intro} \quad \quad \quad [\neg(\varphi \vee \neg\varphi)]^1 \\
\vdots \\
\vdots \\
\varphi \neg\text{Elim} \\
\hline
2 \frac{\perp}{\neg\varphi} \neg\text{Intro} \quad \quad \quad \vdots \\
\vdots \\
\varphi \neg\text{Elim} \\
\hline
1 \frac{\perp}{\varphi \vee \neg\varphi} \perp_C
\end{array}$$

On the right side we use the same strategy, except we get φ by \perp_C :

$$\begin{array}{c}
[\neg(\varphi \vee \neg\varphi)]^1 \quad \frac{[\varphi]^2}{\varphi \vee \neg\varphi} \vee\text{Intro} \quad \quad \quad [\neg(\varphi \vee \neg\varphi)]^1 \quad \frac{[\neg\varphi]^3}{\varphi \vee \neg\varphi} \vee\text{Intro} \\
\vdots \\
\vdots \\
\varphi \neg\text{Elim} \quad \quad \quad \vdots \\
\vdots \\
\varphi \neg\text{Elim} \\
\hline
2 \frac{\perp}{\neg\varphi} \neg\text{Intro} \quad \quad \quad 3 \frac{\perp}{\varphi} \perp_C \\
\vdots \\
\varphi \neg\text{Elim} \\
\hline
1 \frac{\perp}{\varphi \vee \neg\varphi} \perp_C
\end{array}$$

Problem ntd.1. Give **derivations** that show the following:

1. $\varphi \wedge (\psi \wedge \chi) \vdash (\varphi \wedge \psi) \wedge \chi$.
2. $\varphi \vee (\psi \vee \chi) \vdash (\varphi \vee \psi) \vee \chi$.
3. $\varphi \rightarrow (\psi \rightarrow \chi) \vdash \psi \rightarrow (\varphi \rightarrow \chi)$.
4. $\varphi \vdash \neg\neg\varphi$.

Problem ntd.2. Give **derivations** that show the following:

1. $(\varphi \vee \psi) \rightarrow \chi \vdash \varphi \rightarrow \chi$.
2. $(\varphi \rightarrow \chi) \wedge (\psi \rightarrow \chi) \vdash (\varphi \vee \psi) \rightarrow \chi$.
3. $\vdash \neg(\varphi \wedge \neg\varphi)$.
4. $\psi \rightarrow \varphi \vdash \neg\varphi \rightarrow \neg\psi$.
5. $\vdash (\varphi \rightarrow \neg\varphi) \rightarrow \neg\varphi$.
6. $\vdash \neg(\varphi \rightarrow \psi) \rightarrow \neg\psi$.

7. $\varphi \rightarrow \chi \vdash \neg(\varphi \wedge \neg\chi)$.
8. $\varphi \wedge \neg\chi \vdash \neg(\varphi \rightarrow \chi)$.
9. $\varphi \vee \psi, \neg\psi \vdash \varphi$.
10. $\neg\varphi \vee \neg\psi \vdash \neg(\varphi \wedge \psi)$.
11. $\vdash (\neg\varphi \wedge \neg\psi) \rightarrow \neg(\varphi \vee \psi)$.
12. $\vdash \neg(\varphi \vee \psi) \rightarrow (\neg\varphi \wedge \neg\psi)$.

Problem ntd.3. Give **derivations** that show the following:

1. $\neg(\varphi \rightarrow \psi) \vdash \varphi$.
2. $\neg(\varphi \wedge \psi) \vdash \neg\varphi \vee \neg\psi$.
3. $\varphi \rightarrow \psi \vdash \neg\varphi \vee \psi$.
4. $\vdash \neg\neg\varphi \rightarrow \varphi$.
5. $\varphi \rightarrow \psi, \neg\varphi \rightarrow \psi \vdash \psi$.
6. $(\varphi \wedge \psi) \rightarrow \chi \vdash (\varphi \rightarrow \chi) \vee (\psi \rightarrow \chi)$.
7. $(\varphi \rightarrow \psi) \rightarrow \varphi \vdash \varphi$.
8. $\vdash (\varphi \rightarrow \psi) \vee (\psi \rightarrow \chi)$.

(These all require the \perp_C rule.)

ntd.6 Derivations with Quantifiers

Example ntd.7. When dealing with quantifiers, we have to make sure not to violate the eigenvariable condition, and sometimes this requires us to play around with the order of carrying out certain inferences. In general, it helps to try and take care of rules subject to the eigenvariable condition first (they will be lower down in the finished proof).

Let's see how we'd give a **derivation** of the **formula** $\exists x \neg\varphi(x) \rightarrow \neg\forall x \varphi(x)$. Starting as usual, we write

$$\overline{\exists x \neg\varphi(x) \rightarrow \neg\forall x \varphi(x)}$$

We start by writing down what it would take to justify that last step using the \rightarrow Intro rule.

$$\begin{array}{c}
[\exists x \neg\varphi(x)]^1 \\
\vdots \\
\vdots \\
\neg\forall x \varphi(x) \\
\hline
1 \frac{\exists x \neg\varphi(x) \rightarrow \neg\forall x \varphi(x)}{\rightarrow\text{Intro}}
\end{array}$$

Since there is no obvious rule to apply to $\neg\forall x \varphi(x)$, we will proceed by setting up the **derivation** so we can use the $\exists\text{Elim}$ rule. Here we must pay attention to the eigenvariable condition, and choose a constant that does not appear in $\exists x \varphi(x)$ or any assumptions that it depends on. (Since no **constant symbols** appear, however, any choice will do fine.)

$$\begin{array}{c}
[\neg\varphi(a)]^2 \\
\vdots \\
\vdots \\
\frac{[\exists x \neg\varphi(x)]^1 \quad \neg\forall x \varphi(x)}{\exists\text{Elim}} \\
\hline
1 \frac{\exists x \neg\varphi(x) \rightarrow \neg\forall x \varphi(x)}{\rightarrow\text{Intro}}
\end{array}$$

In order to derive $\neg\forall x \varphi(x)$, we will attempt to use the $\neg\text{Intro}$ rule: this requires that we derive a contradiction, possibly using $\forall x \varphi(x)$ as an additional assumption. Of course, this contradiction may involve the assumption $\neg\varphi(a)$ which will be discharged by the $\exists\text{Elim}$ inference. We can set it up as follows:

$$\begin{array}{c}
[\neg\varphi(a)]^2, [\forall x \varphi(x)]^3 \\
\vdots \\
\vdots \\
\perp \\
\hline
3 \frac{\perp}{\neg\forall x \varphi(x)} \neg\text{Intro} \\
\hline
2 \frac{[\exists x \neg\varphi(x)]^1 \quad \neg\forall x \varphi(x)}{\exists\text{Elim}} \\
\hline
1 \frac{\exists x \neg\varphi(x) \rightarrow \neg\forall x \varphi(x)}{\rightarrow\text{Intro}}
\end{array}$$

It looks like we are close to getting a contradiction. The easiest rule to apply is the $\forall\text{Elim}$, which has no eigenvariable conditions. Since we can use any term we want to replace the universally quantified x , it makes the most sense to continue using a so we can reach a contradiction.

$$\begin{array}{c}
\frac{[\forall x \varphi(x)]^3}{\varphi(a)} \forall\text{Elim} \\
\hline
[\neg\varphi(a)]^2 \quad \varphi(a) \\
\hline
\neg\text{Elim} \\
\hline
3 \frac{\perp}{\neg\forall x \varphi(x)} \neg\text{Intro} \\
\hline
2 \frac{[\exists x \neg\varphi(x)]^1 \quad \neg\forall x \varphi(x)}{\exists\text{Elim}} \\
\hline
1 \frac{\exists x \neg\varphi(x) \rightarrow \neg\forall x \varphi(x)}{\rightarrow\text{Intro}}
\end{array}$$

It is important, especially when dealing with quantifiers, to double check at this point that the eigenvariable condition has not been violated. Since the only rule we applied that is subject to the eigenvariable condition was \exists Elim, and the eigenvariable a does not occur in any assumptions it depends on, this is a correct **derivation**.

Example ntd.8. Sometimes we may derive a **formula** from other **formulas**. In these cases, we may have undischarged assumptions. It is important to keep track of our assumptions as well as the end goal.

Let's see how we'd give a **derivation** of the **formula** $\exists x \chi(x, b)$ from the assumptions $\exists x (\varphi(x) \wedge \psi(x))$ and $\forall x (\psi(x) \rightarrow \chi(x, b))$. Starting as usual, we write the conclusion at the bottom.

$$\overline{\exists x \chi(x, b)}$$

We have two premises to work with. To use the first, i.e., try to find a **derivation** of $\exists x \chi(x, b)$ from $\exists x (\varphi(x) \wedge \psi(x))$ we would use the \exists Elim rule. Since it has an eigenvariable condition, we will apply that rule first. We get the following:

$$\begin{array}{c}
 [\varphi(a) \wedge \psi(a)]^1 \\
 \vdots \\
 \vdots \\
 \vdots \\
 \frac{1 \quad \exists x (\varphi(x) \wedge \psi(x)) \quad \exists x \chi(x, b)}{\exists x \chi(x, b)} \exists\text{Elim}
 \end{array}$$

The two assumptions we are working with share ψ . It may be useful at this point to apply \wedge Elim to separate out $\psi(a)$.

$$\begin{array}{c}
 \frac{[\varphi(a) \wedge \psi(a)]^1}{\psi(a)} \wedge\text{Elim} \\
 \vdots \\
 \vdots \\
 \vdots \\
 \frac{1 \quad \exists x (\varphi(x) \wedge \psi(x)) \quad \exists x \chi(x, b)}{\exists x \chi(x, b)} \exists\text{Elim}
 \end{array}$$

The second assumption we have to work with is $\forall x (\psi(x) \rightarrow \chi(x, b))$. Since there is no eigenvariable condition we can instantiate x with the **constant symbol** a using \forall Elim to get $\psi(a) \rightarrow \chi(a, b)$. We now have both $\psi(a) \rightarrow \chi(a, b)$ and $\psi(a)$. Our next move should be a straightforward application of the \rightarrow Elim rule.

$$\begin{array}{c}
\frac{\frac{\forall x (\psi(x) \rightarrow \chi(x, b))}{\psi(a) \rightarrow \chi(a, b)} \forall\text{Elim} \quad \frac{[\varphi(a) \wedge \psi(a)]^1}{\psi(a)} \wedge\text{Elim}}{\chi(a, b)} \rightarrow\text{Elim} \\
\vdots \\
\frac{\exists x (\varphi(x) \wedge \psi(x)) \quad \exists x \chi(x, b)}{\exists x \chi(x, b)} \exists\text{Elim}
\end{array}$$

We are so close! One application of $\exists\text{Intro}$ and we have reached our goal.

$$\begin{array}{c}
\frac{\frac{\forall x (\psi(x) \rightarrow \chi(x, b))}{\psi(a) \rightarrow \chi(a, b)} \forall\text{Elim} \quad \frac{[\varphi(a) \wedge \psi(a)]^1}{\psi(a)} \wedge\text{Elim}}{\chi(a, b)} \rightarrow\text{Elim} \\
\frac{\exists x (\varphi(x) \wedge \psi(x)) \quad \frac{\chi(a, b)}{\exists x \chi(x, b)} \exists\text{Intro}}{\exists x \chi(x, b)} \exists\text{Elim}
\end{array}$$

Since we ensured at each step that the eigenvariable conditions were not violated, we can be confident that this is a correct **derivation**.

Example ntd.9. Give a **derivation** of the **formula** $\neg\forall x \varphi(x)$ from the assumptions $\forall x \varphi(x) \rightarrow \exists y \psi(y)$ and $\neg\exists y \psi(y)$. Starting as usual, we write the target **formula** at the bottom.

$$\overline{\neg\forall x \varphi(x)}$$

The last line of the **derivation** is a negation, so let's try using $\neg\text{Intro}$. This will require that we figure out how to **derive** a contradiction.

$$\begin{array}{c}
[\forall x \varphi(x)]^1 \\
\vdots \\
\vdots \\
\perp \\
\frac{}{\neg\forall x \varphi(x)} \neg\text{Intro}
\end{array}$$

So far so good. We can use $\forall\text{Elim}$ but it's not obvious if that will help us get to our goal. Instead, let's use one of our assumptions. $\forall x \varphi(x) \rightarrow \exists y \psi(y)$ together with $\forall x \varphi(x)$ will allow us to use the $\rightarrow\text{Elim}$ rule.

$$\begin{array}{c}
\frac{\forall x \varphi(x) \rightarrow \exists y \psi(y) \quad [\forall x \varphi(x)]^1}{\exists y \psi(y)} \rightarrow\text{Elim} \\
\vdots \\
\vdots \\
\perp \\
\frac{}{\neg\forall x \varphi(x)} \neg\text{Intro}
\end{array}$$

We now have one final assumption to work with, and it looks like this will help us reach a contradiction by using \neg Elim.

$$\frac{\frac{\neg\exists y \psi(y) \quad \frac{\forall x \varphi(x) \rightarrow \exists y \psi(y) \quad [\forall x \varphi(x)]^1}{\exists y \psi(y)} \rightarrow\text{Elim}}{1 \quad \frac{\perp}{\neg\forall x \varphi(x)} \neg\text{Intro}}{\neg\text{Elim}}$$

Problem ntd.4. Give **derivations** that show the following:

1. $\vdash (\forall x \varphi(x) \wedge \forall y \psi(y)) \rightarrow \forall z (\varphi(z) \wedge \psi(z))$.
2. $\vdash (\exists x \varphi(x) \vee \exists y \psi(y)) \rightarrow \exists z (\varphi(z) \vee \psi(z))$.
3. $\forall x (\varphi(x) \rightarrow \psi) \vdash \exists y \varphi(y) \rightarrow \psi$.
4. $\forall x \neg\varphi(x) \vdash \neg\exists x \varphi(x)$.
5. $\vdash \neg\exists x \varphi(x) \rightarrow \forall x \neg\varphi(x)$.
6. $\vdash \neg\exists x \forall y ((\varphi(x, y) \rightarrow \neg\varphi(y, y)) \wedge (\neg\varphi(y, y) \rightarrow \varphi(x, y)))$.

Problem ntd.5. Give **derivations** that show the following:

1. $\vdash \neg\forall x \varphi(x) \rightarrow \exists x \neg\varphi(x)$.
2. $(\forall x \varphi(x) \rightarrow \psi) \vdash \exists y (\varphi(y) \rightarrow \psi)$.
3. $\vdash \exists x (\varphi(x) \rightarrow \forall y \varphi(y))$.

(These all require the \perp_C rule.)

ntd.7 Proof-Theoretic Notions

fol:ntd:ptn:
sec

This section collects the definitions the provability relation and consistency for natural deduction.

explanation Just as we've defined a number of important semantic notions (validity, entailment, satisfiability), we now define corresponding *proof-theoretic notions*. These are not defined by appeal to satisfaction of **sentences** in **structures**, but by appeal to the **derivability** or **non-derivability** of certain **sentences** from others. It was an important discovery that these notions coincide. That they do is the content of the *soundness* and *completeness theorems*.

Definition ntd.10 (Theorems). A **sentence** φ is a *theorem* if there is a **derivation** of φ in natural deduction in which all assumptions are **discharged**. We write $\vdash \varphi$ if φ is a theorem and $\not\vdash \varphi$ if it is not.

Definition ntd.11 (Derivability). A sentence φ is *derivable* from a set of sentences Γ , $\Gamma \vdash \varphi$, if there is a *derivation* with conclusion φ and in which every assumption is either *discharged* or is in Γ . If φ is not *derivable* from Γ we write $\Gamma \not\vdash \varphi$.

Definition ntd.12 (Consistency). A set of sentences Γ is *inconsistent* iff $\Gamma \vdash \perp$. If Γ is not inconsistent, i.e., if $\Gamma \not\vdash \perp$, we say it is *consistent*.

*fol:ntd:ptn:
prop:reflexivity*

Proposition ntd.13 (Reflexivity). If $\varphi \in \Gamma$, then $\Gamma \vdash \varphi$.

Proof. The assumption φ by itself is a *derivation* of φ where every *undischarged* assumption (i.e., φ) is in Γ . \square

*fol:ntd:ptn:
prop:monotonicity*

Proposition ntd.14 (Monotonicity). If $\Gamma \subseteq \Delta$ and $\Gamma \vdash \varphi$, then $\Delta \vdash \varphi$.

Proof. Any *derivation* of φ from Γ is also a *derivation* of φ from Δ . \square

*fol:ntd:ptn:
prop:transitivity*

Proposition ntd.15 (Transitivity). If $\Gamma \vdash \varphi$ and $\{\varphi\} \cup \Delta \vdash \psi$, then $\Gamma \cup \Delta \vdash \psi$.

Proof. If $\Gamma \vdash \varphi$, there is a *derivation* δ_0 of φ with all *undischarged* assumptions in Γ . If $\{\varphi\} \cup \Delta \vdash \psi$, then there is a *derivation* δ_1 of ψ with all *undischarged* assumptions in $\{\varphi\} \cup \Delta$. Now consider:

$$\begin{array}{ccc}
 \Delta, [\varphi]^1 & & \Gamma \\
 \vdots & & \vdots \\
 \delta_1 & & \delta_0 \\
 \vdots & & \vdots \\
 \psi & & \varphi \\
 \hline
 \varphi \rightarrow \psi & \xrightarrow{\text{Intro}} & \varphi \\
 \hline
 \psi & & \psi \\
 & & \xrightarrow{\text{Elim}}
 \end{array}$$

The *undischarged* assumptions are now all among $\Gamma \cup \Delta$, so this shows $\Gamma \cup \Delta \vdash \psi$. \square

When $\Gamma = \{\varphi_1, \varphi_2, \dots, \varphi_k\}$ is a finite set we may use the simplified notation $\varphi_1, \varphi_2, \dots, \varphi_k \vdash \psi$ for $\Gamma \vdash \psi$, in particular $\varphi \vdash \psi$ means that $\{\varphi\} \vdash \psi$.

Note that if $\Gamma \vdash \varphi$ and $\varphi \vdash \psi$, then $\Gamma \vdash \psi$. It follows also that if $\varphi_1, \dots, \varphi_n \vdash \psi$ and $\Gamma \vdash \varphi_i$ for each i , then $\Gamma \vdash \psi$.

*fol:ntd:ptn:
prop:incons*

Proposition ntd.16. The following are equivalent.

1. Γ is inconsistent.
2. $\Gamma \vdash \varphi$ for every *sentence* φ .
3. $\Gamma \vdash \varphi$ and $\Gamma \vdash \neg\varphi$ for some *sentence* φ .

Proof. Exercise. \square

Problem ntd.6. Prove [Proposition ntd.16](#)

Proposition ntd.17 (Compactness).

*fol:ntd:ptn:
prop:proves-compact*

1. If $\Gamma \vdash \varphi$ then there is a finite subset $\Gamma_0 \subseteq \Gamma$ such that $\Gamma_0 \vdash \varphi$.
2. If every finite subset of Γ is consistent, then Γ is consistent.

Proof. 1. If $\Gamma \vdash \varphi$, then there is a **derivation** δ of φ from Γ . Let Γ_0 be the set of **undischarged** assumptions of δ . Since any **derivation** is finite, Γ_0 can only contain finitely many **sentences**. So, δ is a **derivation** of φ from a finite $\Gamma_0 \subseteq \Gamma$.

2. This is the contrapositive of (1) for the special case $\varphi \equiv \perp$. □

ntd.8 Derivability and Consistency

We will now establish a number of properties of the **derivability** relation. They are independently interesting, but each will play a role in the proof of the completeness theorem.

*fol:ntd:prv:
sec*

Proposition ntd.18. *If $\Gamma \vdash \varphi$ and $\Gamma \cup \{\varphi\}$ is inconsistent, then Γ is inconsistent.*

*fol:ntd:prv:
prop:provability-contr*

Proof. Let the **derivation** of φ from Γ be δ_1 and the **derivation** of \perp from $\Gamma \cup \{\varphi\}$ be δ_2 . We can then **derive**:

$$\frac{\begin{array}{c} \Gamma, [\varphi]^1 \\ \vdots \\ \delta_2 \\ \vdots \\ \perp \\ \hline \neg\varphi \quad \neg\text{Intro} \end{array}}{\perp} \quad \frac{\begin{array}{c} \Gamma \\ \vdots \\ \delta_1 \\ \vdots \\ \varphi \\ \hline \neg\text{Elim} \end{array}}{\perp} \quad \neg\text{Elim}$$

In the new **derivation**, the assumption φ is **discharged**, so it is a **derivation** from Γ . □

Proposition ntd.19. *$\Gamma \vdash \varphi$ iff $\Gamma \cup \{\neg\varphi\}$ is inconsistent.*

*fol:ntd:prv:
prop:prov-incons*

Proof. First suppose $\Gamma \vdash \varphi$, i.e., there is a **derivation** δ_0 of φ from **undischarged** assumptions Γ . We obtain a **derivation** of \perp from $\Gamma \cup \{\neg\varphi\}$ as follows:

$$\frac{\begin{array}{c} \Gamma \\ \vdots \\ \delta_0 \\ \vdots \\ \varphi \\ \hline \neg\varphi \quad \neg\text{Elim} \end{array}}{\perp} \quad \neg\text{Elim}$$

Now assume $\Gamma \cup \{\neg\varphi\}$ is inconsistent, and let δ_1 be the corresponding derivation of \perp from undischarged assumptions in $\Gamma \cup \{\neg\varphi\}$. We obtain a derivation of φ from Γ alone by using \perp_C :

$$\frac{\Gamma, [\neg\varphi]^1 \quad \begin{array}{c} \vdots \\ \delta_1 \\ \vdots \\ \perp \end{array}}{1 \frac{\perp}{\varphi} \perp_C} \quad \square$$

Problem ntd.7. Prove that $\Gamma \vdash \neg\varphi$ iff $\Gamma \cup \{\varphi\}$ is inconsistent.

*fol.ntd.prv:
prop:explicit-inc*

Proposition ntd.20. *If $\Gamma \vdash \varphi$ and $\neg\varphi \in \Gamma$, then Γ is inconsistent.*

Proof. Suppose $\Gamma \vdash \varphi$ and $\neg\varphi \in \Gamma$. Then there is a derivation δ of φ from Γ . Consider this simple application of the \neg -Elim rule:

$$\frac{\Gamma \quad \begin{array}{c} \vdots \\ \delta \\ \vdots \\ \varphi \end{array} \quad \neg\varphi}{\perp} \neg\text{Elim}$$

Since $\neg\varphi \in \Gamma$, all undischarged assumptions are in Γ , this shows that $\Gamma \vdash \perp$. \square

*fol.ntd.prv:
prop:provability-exhaustive*

Proposition ntd.21. *If $\Gamma \cup \{\varphi\}$ and $\Gamma \cup \{\neg\varphi\}$ are both inconsistent, then Γ is inconsistent.*

Proof. There are derivations δ_1 and δ_2 of \perp from $\Gamma \cup \{\varphi\}$ and \perp from $\Gamma \cup \{\neg\varphi\}$, respectively. We can then derive

$$\frac{\Gamma, [\neg\varphi]^2 \quad \begin{array}{c} \vdots \\ \delta_2 \\ \vdots \\ \perp \end{array} \quad \Gamma, [\varphi]^1 \quad \begin{array}{c} \vdots \\ \delta_1 \\ \vdots \\ \perp \end{array}}{2 \frac{\perp}{\neg\neg\varphi} \neg\text{Intro} \quad 1 \frac{\perp}{\neg\varphi} \neg\text{Intro} \quad \perp} \neg\text{Elim}$$

Since the assumptions φ and $\neg\varphi$ are discharged, this is a derivation of \perp from Γ alone. Hence Γ is inconsistent. \square

ntd.9 Derivability and the Propositional Connectives

explanation We establish that the **derivability** relation \vdash of natural deduction is strong enough to establish some basic facts involving the propositional connectives, such as that $\varphi \wedge \psi \vdash \varphi$ and $\varphi, \varphi \rightarrow \psi \vdash \psi$ (modus ponens). These facts are needed for the proof of the completeness theorem.

fol:ntd:ppr:
sec

Proposition ntd.22.

1. Both $\varphi \wedge \psi \vdash \varphi$ and $\varphi \wedge \psi \vdash \psi$
2. $\varphi, \psi \vdash \varphi \wedge \psi$.

fol:ntd:ppr:
prop:provability-land
fol:ntd:ppr:
prop:provability-land-left
fol:ntd:ppr:
prop:provability-land-right

Proof. 1. We can **derive** both

$$\frac{\varphi \wedge \psi}{\varphi} \wedge\text{Elim} \qquad \frac{\varphi \wedge \psi}{\psi} \wedge\text{Elim}$$

2. We can **derive**:

$$\frac{\varphi \quad \psi}{\varphi \wedge \psi} \wedge\text{Intro}$$

□

Proposition ntd.23.

1. $\varphi \vee \psi, \neg\varphi, \neg\psi$ is inconsistent.
2. Both $\varphi \vdash \varphi \vee \psi$ and $\psi \vdash \varphi \vee \psi$.

fol:ntd:ppr:
prop:provability-lor

Proof. 1. Consider the following **derivation**:

$$\frac{1 \quad \varphi \vee \psi \quad \frac{\frac{\neg\varphi \quad [\varphi]^1}{\perp} \neg\text{Elim} \quad \frac{\neg\psi \quad [\psi]^1}{\perp} \neg\text{Elim}}{\perp} \vee\text{Elim}}{\perp}$$

This is a **derivation** of \perp from **undischarged** assumptions $\varphi \vee \psi$, $\neg\varphi$, and $\neg\psi$.

2. We can **derive** both

$$\frac{\varphi}{\varphi \vee \psi} \vee\text{Intro} \qquad \frac{\psi}{\varphi \vee \psi} \vee\text{Intro}$$

□

Proposition ntd.24.

1. $\varphi, \varphi \rightarrow \psi \vdash \psi$.
2. Both $\neg\varphi \vdash \varphi \rightarrow \psi$ and $\psi \vdash \varphi \rightarrow \psi$.

fol:ntd:ppr:
prop:provability-lif
fol:ntd:ppr:
prop:provability-lif-left
fol:ntd:ppr:
prop:provability-lif-right

Proof. 1. We can **derive**:

$$\frac{\varphi \rightarrow \psi \quad \varphi}{\psi} \rightarrow\text{Elim}$$

2. This is shown by the following two **derivations**:

$$\frac{\frac{\frac{\perp}{\psi} \perp_I}{\varphi \rightarrow \psi} \rightarrow\text{Intro} \quad \frac{\neg\varphi \quad [\varphi]^1}{\perp} \neg\text{Elim}}{\varphi \rightarrow \psi} \rightarrow\text{Intro} \quad \frac{\psi}{\varphi \rightarrow \psi} \rightarrow\text{Intro}$$

Note that $\rightarrow\text{Intro}$ may, but does not have to, **discharge** the assumption φ .
□

ntd.10 Derivability and the Quantifiers

fol:ntd:qpr:sec The completeness theorem also requires that the natural deduction rules yield explanation the facts about \vdash established in this section.

fol:ntd:qpr:thm:strong-generalization **Theorem ntd.25.** *If c is a constant not occurring in Γ or $\varphi(x)$ and $\Gamma \vdash \varphi(c)$, then $\Gamma \vdash \forall x \varphi(x)$.*

Proof. Let δ be a **derivation** of $\varphi(c)$ from Γ . By adding a $\forall\text{Intro}$ inference, we obtain a **derivation** of $\forall x \varphi(x)$. Since c does not occur in Γ or $\varphi(x)$, the eigenvariable condition is satisfied. □

fol:ntd:qpr:prop:provability-quantifiers **Proposition ntd.26.**

1. $\varphi(t) \vdash \exists x \varphi(x)$.
2. $\forall x \varphi(x) \vdash \varphi(t)$.

Proof. 1. The following is a **derivation** of $\exists x \varphi(x)$ from $\varphi(t)$:

$$\frac{\varphi(t)}{\exists x \varphi(x)} \exists\text{Intro}$$

2. The following is a **derivation** of $\varphi(t)$ from $\forall x \varphi(x)$:

$$\frac{\forall x \varphi(x)}{\varphi(t)} \forall\text{Elim} \quad \square$$

ntd.11 Soundness

explanation A **derivation** system, such as natural deduction, is *sound* if it cannot **derive** things that do not actually follow. Soundness is thus a kind of guaranteed safety property for **derivation** systems. Depending on which proof theoretic property is in question, we would like to know for instance, that fol:ntd:sou:sec

1. every **derivable sentence** is valid;
2. if a **sentence** is **derivable** from some others, it is also a consequence of them;
3. if a set of **sentences** is inconsistent, it is unsatisfiable.

These are important properties of a **derivation** system. If any of them do not hold, the **derivation** system is deficient—it would **derive** too much. Consequently, establishing the soundness of a **derivation** system is of the utmost importance.

Theorem ntd.27 (Soundness). *If φ is **derivable** from the **undischarged** assumptions Γ , then $\Gamma \vDash \varphi$.* fol:ntd:sou:thm:soundness

Proof. Let δ be a **derivation** of φ . We proceed by induction on the number of inferences in δ .

For the induction basis we show the claim if the number of inferences is 0. In this case, δ consists only of a single **sentence** φ , i.e., an assumption. That assumption is **undischarged**, since assumptions can only be **discharged** by inferences, and there are no inferences. So, any **structure** \mathfrak{M} that satisfies all of the **undischarged** assumptions of the proof also satisfies φ .

Now for the inductive step. Suppose that δ contains n inferences. The premise(s) of the lowermost inference are **derived** using sub-**derivations**, each of which contains fewer than n inferences. We assume the induction hypothesis: The premises of the lowermost inference follow from the **undischarged** assumptions of the sub-**derivations** ending in those premises. We have to show that the conclusion φ follows from the **undischarged** assumptions of the entire proof.

We distinguish cases according to the type of the lowermost inference. First, we consider the possible inferences with only one premise.

1. Suppose that the last inference is \neg -Intro: The **derivation** has the form

$$\frac{\Gamma, [\varphi]^n \quad \begin{array}{c} \vdots \\ \delta_1 \\ \vdots \end{array}}{n \frac{\perp}{\neg\varphi} \neg\text{Intro}}$$

By inductive hypothesis, \perp follows from the **undischarged** assumptions $\Gamma \cup \{\varphi\}$ of δ_1 . Consider a **structure** \mathfrak{M} . We need to show that, if $\mathfrak{M} \models \Gamma$, then $\mathfrak{M} \models \neg\varphi$. Suppose for reductio that $\mathfrak{M} \models \Gamma$, but $\mathfrak{M} \not\models \neg\varphi$, i.e., $\mathfrak{M} \models \varphi$. This would mean that $\mathfrak{M} \models \Gamma \cup \{\varphi\}$. This is contrary to our inductive hypothesis. So, $\mathfrak{M} \models \neg\varphi$.

2. The last inference is \wedge Elim: There are two variants: φ or ψ may be inferred from the premise $\varphi \wedge \psi$. Consider the first case. The **derivation** δ looks like this:

$$\frac{\begin{array}{c} \Gamma \\ \vdots \\ \delta_1 \\ \vdots \\ \varphi \wedge \psi \end{array}}{\varphi} \wedge\text{Elim}$$

By inductive hypothesis, $\varphi \wedge \psi$ follows from the **undischarged** assumptions Γ of δ_1 . Consider a **structure** \mathfrak{M} . We need to show that, if $\mathfrak{M} \models \Gamma$, then $\mathfrak{M} \models \varphi$. Suppose $\mathfrak{M} \models \Gamma$. By our inductive hypothesis ($\Gamma \models \varphi \wedge \psi$), we know that $\mathfrak{M} \models \varphi \wedge \psi$. By definition, $\mathfrak{M} \models \varphi \wedge \psi$ iff $\mathfrak{M} \models \varphi$ and $\mathfrak{M} \models \psi$. (The case where ψ is inferred from $\varphi \wedge \psi$ is handled similarly.)

3. The last inference is \vee Intro: There are two variants: $\varphi \vee \psi$ may be inferred from the premise φ or the premise ψ . Consider the first case. The **derivation** has the form

$$\frac{\begin{array}{c} \Gamma \\ \vdots \\ \delta_1 \\ \vdots \\ \varphi \end{array}}{\varphi \vee \psi} \vee\text{Intro}$$

By inductive hypothesis, φ follows from the **undischarged** assumptions Γ of δ_1 . Consider a **structure** \mathfrak{M} . We need to show that, if $\mathfrak{M} \models \Gamma$, then $\mathfrak{M} \models \varphi \vee \psi$. Suppose $\mathfrak{M} \models \Gamma$; then $\mathfrak{M} \models \varphi$ since $\Gamma \models \varphi$ (the inductive hypothesis). So it must also be the case that $\mathfrak{M} \models \varphi \vee \psi$. (The case where $\varphi \vee \psi$ is inferred from ψ is handled similarly.)

4. The last inference is \rightarrow Intro: $\varphi \rightarrow \psi$ is inferred from a subproof with assumption φ and conclusion ψ , i.e.,

$${}^n \frac{\begin{array}{c} \Gamma, [\varphi]^n \\ \vdots \\ \delta_1 \\ \vdots \\ \psi \end{array}}{\varphi \rightarrow \psi} \rightarrow\text{Intro}$$

By inductive hypothesis, ψ follows from the **undischarged** assumptions of δ_1 , i.e., $\Gamma \cup \{\varphi\} \vDash \psi$. Consider a **structure** \mathfrak{M} . The **undischarged** assumptions of δ are just Γ , since φ is discharged at the last inference. So we need to show that $\Gamma \vDash \varphi \rightarrow \psi$. For reductio, suppose that for some **structure** \mathfrak{M} , $\mathfrak{M} \vDash \Gamma$ but $\mathfrak{M} \not\vDash \varphi \rightarrow \psi$. So, $\mathfrak{M} \vDash \varphi$ and $\mathfrak{M} \not\vDash \psi$. But by hypothesis, ψ is a consequence of $\Gamma \cup \{\varphi\}$, i.e., $\mathfrak{M} \vDash \psi$, which is a contradiction. So, $\Gamma \vDash \varphi \rightarrow \psi$.

5. The last inference is \perp_I : Here, δ ends in

$$\frac{\begin{array}{c} \Gamma \\ \vdots \\ \delta_1 \\ \vdots \\ \perp \end{array}}{\varphi} \perp_I$$

By induction hypothesis, $\Gamma \vDash \perp$. We have to show that $\Gamma \vDash \varphi$. Suppose not; then for some \mathfrak{M} we have $\mathfrak{M} \vDash \Gamma$ and $\mathfrak{M} \not\vDash \varphi$. But we always have $\mathfrak{M} \not\vDash \perp$, so this would mean that $\Gamma \not\vDash \perp$, contrary to the induction hypothesis.

6. The last inference is \perp_C : Exercise.
 7. The last inference is \forall Intro: Then δ has the form

$$\frac{\begin{array}{c} \Gamma \\ \vdots \\ \delta_1 \\ \vdots \\ \varphi(a) \end{array}}{\forall x \varphi(x)} \forall\text{Intro}$$

The premise $\varphi(a)$ is a consequence of the **undischarged** assumptions Γ by induction hypothesis. Consider some structure, \mathfrak{M} , such that $\mathfrak{M} \vDash \Gamma$. We need to show that $\mathfrak{M} \vDash \forall x \varphi(x)$. Since $\forall x \varphi(x)$ is a **sentence**, this means we have to show that for every variable assignment s , $\mathfrak{M}, s \vDash \varphi(x)$ (??). Since Γ consists entirely of sentences, $\mathfrak{M}, s \vDash \psi$ for all $\psi \in \Gamma$ by ???. Let \mathfrak{M}' be like \mathfrak{M} except that $a^{\mathfrak{M}'} = s(x)$. Since a does not occur in Γ , $\mathfrak{M}' \vDash \Gamma$ by ???. Since $\Gamma \vDash \varphi(a)$, $\mathfrak{M}' \vDash \varphi(a)$. Since $\varphi(a)$ is a **sentence**, $\mathfrak{M}', s \vDash \varphi(a)$ by ???. $\mathfrak{M}', s \vDash \varphi(x)$ iff $\mathfrak{M}' \vDash \varphi(a)$ by ?? (recall that $\varphi(a)$ is just $\varphi(x)[a/x]$). So, $\mathfrak{M}', s \vDash \varphi(x)$. Since a does not occur in $\varphi(x)$, by ??, $\mathfrak{M}, s \vDash \varphi(x)$. But s was an arbitrary variable assignment, so $\mathfrak{M} \vDash \forall x \varphi(x)$.

8. The last inference is \exists Intro: Exercise.
 9. The last inference is \forall Elim: Exercise.

Now let's consider the possible inferences with several premises: \forall Elim, \wedge Intro, \rightarrow Elim, and \exists Elim.

1. The last inference is \wedge Intro. $\varphi \wedge \psi$ is inferred from the premises φ and ψ and δ has the form

$$\frac{\begin{array}{c} \Gamma_1 \\ \vdots \\ \delta_1 \\ \vdots \\ \varphi \end{array} \quad \begin{array}{c} \Gamma_2 \\ \vdots \\ \delta_2 \\ \vdots \\ \psi \end{array}}{\varphi \wedge \psi} \wedge\text{Intro}$$

By induction hypothesis, φ follows from the **undischarged** assumptions Γ_1 of δ_1 and ψ follows from the **undischarged** assumptions Γ_2 of δ_2 . The **undischarged** assumptions of δ are $\Gamma_1 \cup \Gamma_2$, so we have to show that $\Gamma_1 \cup \Gamma_2 \models \varphi \wedge \psi$. Consider a **structure** \mathfrak{M} with $\mathfrak{M} \models \Gamma_1 \cup \Gamma_2$. Since $\mathfrak{M} \models \Gamma_1$, it must be the case that $\mathfrak{M} \models \varphi$ as $\Gamma_1 \models \varphi$, and since $\mathfrak{M} \models \Gamma_2$, $\mathfrak{M} \models \psi$ since $\Gamma_2 \models \psi$. Together, $\mathfrak{M} \models \varphi \wedge \psi$.

2. The last inference is \forall Elim: Exercise.
3. The last inference is \rightarrow Elim. ψ is inferred from the premises $\varphi \rightarrow \psi$ and φ . The **derivation** δ looks like this:

$$\frac{\begin{array}{c} \Gamma_1 \\ \vdots \\ \delta_1 \\ \vdots \\ \varphi \rightarrow \psi \end{array} \quad \begin{array}{c} \Gamma_2 \\ \vdots \\ \delta_2 \\ \vdots \\ \varphi \end{array}}{\psi} \rightarrow\text{Elim}$$

By induction hypothesis, $\varphi \rightarrow \psi$ follows from the **undischarged** assumptions Γ_1 of δ_1 and φ follows from the **undischarged** assumptions Γ_2 of δ_2 . Consider a **structure** \mathfrak{M} . We need to show that, if $\mathfrak{M} \models \Gamma_1 \cup \Gamma_2$, then $\mathfrak{M} \models \psi$. Suppose $\mathfrak{M} \models \Gamma_1 \cup \Gamma_2$. Since $\Gamma_1 \models \varphi \rightarrow \psi$, $\mathfrak{M} \models \varphi \rightarrow \psi$. Since $\Gamma_2 \models \varphi$, we have $\mathfrak{M} \models \varphi$. This means that $\mathfrak{M} \models \psi$ (For if $\mathfrak{M} \not\models \psi$, since $\mathfrak{M} \models \varphi$, we'd have $\mathfrak{M} \not\models \varphi \rightarrow \psi$, contradicting $\mathfrak{M} \models \varphi \rightarrow \psi$).

4. The last inference is \neg Elim: Exercise.
5. The last inference is \exists Elim: Exercise. □

Problem ntd.8. Complete the proof of **Theorem ntd.27**.

fol:ntd:sou:
cor:weak-soundness **Corollary ntd.28.** *If $\vdash \varphi$, then φ is valid.*

fol:ntd:sou:
cor:consistency-soundness **Corollary ntd.29.** *If Γ is satisfiable, then it is consistent.*

Proof. We prove the contrapositive. Suppose that Γ is not consistent. Then $\Gamma \vdash \perp$, i.e., there is a derivation of \perp from undischarged assumptions in Γ . By Theorem ntd.27, any structure \mathfrak{M} that satisfies Γ must satisfy \perp . Since $\mathfrak{M} \not\models \perp$ for every structure \mathfrak{M} , no \mathfrak{M} can satisfy Γ , i.e., Γ is not satisfiable. \square

ntd.12 Derivations with Identity predicate

Derivations with identity predicate require additional inference rules.

fol:ntd:ide:
sec

$\frac{}{t = t} =\text{Intro}$	$\frac{t_1 = t_2 \quad \varphi(t_1)}{\varphi(t_2)} =\text{Elim}$ $\frac{t_1 = t_2 \quad \varphi(t_2)}{\varphi(t_1)} =\text{Elim}$
--------------------------------	---

In the above rules, t , t_1 , and t_2 are closed terms. The =Intro rule allows us to derive any identity statement of the form $t = t$ outright, from no assumptions.

Example ntd.30. If s and t are closed terms, then $\varphi(s), s = t \vdash \varphi(t)$:

$$\frac{s = t \quad \varphi(s)}{\varphi(t)} =\text{Elim}$$

This may be familiar as the “principle of substitutability of identicals,” or Leibniz’ Law.

Problem ntd.9. Prove that = is both symmetric and transitive, i.e., give derivations of $\forall x \forall y (x = y \rightarrow y = x)$ and $\forall x \forall y \forall z ((x = y \wedge y = z) \rightarrow x = z)$

Example ntd.31. We derive the sentence

$$\forall x \forall y ((\varphi(x) \wedge \varphi(y)) \rightarrow x = y)$$

from the sentence

$$\exists x \forall y (\varphi(y) \rightarrow y = x)$$

We develop the derivation backwards:

$$\begin{array}{c} \exists x \forall y (\varphi(y) \rightarrow y = x) \quad [\varphi(a) \wedge \varphi(b)]^1 \\ \vdots \\ a = b \\ \hline \frac{1 \quad (\varphi(a) \wedge \varphi(b)) \rightarrow a = b}{\forall y ((\varphi(a) \wedge \varphi(y)) \rightarrow a = y)} \rightarrow\text{Intro} \\ \hline \frac{\forall y ((\varphi(a) \wedge \varphi(y)) \rightarrow a = y)}{\forall x \forall y ((\varphi(x) \wedge \varphi(y)) \rightarrow x = y)} \forall\text{Intro} \end{array}$$

We'll now have to use the main assumption: since it is an existential **formula**, we use \exists Elim to **derive** the intermediary conclusion $a = b$.

$$\begin{array}{c}
 [\forall y (\varphi(y) \rightarrow y = c)]^2 \\
 [\varphi(a) \wedge \varphi(b)]^1 \\
 \vdots \\
 \vdots \\
 \vdots \\
 \frac{\exists x \forall y (\varphi(y) \rightarrow y = x) \quad a = b}{a = b} \exists\text{Elim} \\
 \frac{\frac{\frac{a = b}{((\varphi(a) \wedge \varphi(b)) \rightarrow a = b)} \rightarrow\text{Intro}}{\forall y ((\varphi(a) \wedge \varphi(y)) \rightarrow a = y)} \forall\text{Intro}}{\forall x \forall y ((\varphi(x) \wedge \varphi(y)) \rightarrow x = y)} \forall\text{Intro}
 \end{array}$$

The sub-**derivation** on the top right is completed by using its assumptions to show that $a = c$ and $b = c$. This requires two separate **derivations**. The **derivation** for $a = c$ is as follows:

$$\frac{\frac{[\forall y (\varphi(y) \rightarrow y = c)]^2}{\varphi(a) \rightarrow a = c} \forall\text{Elim} \quad \frac{[\varphi(a) \wedge \varphi(b)]^1}{\varphi(a)} \wedge\text{Elim}}{a = c} \rightarrow\text{Elim}$$

From $a = c$ and $b = c$ we **derive** $a = b$ by $=$ Elim.

Problem ntd.10. Give **derivations** of the following **formulas**:

1. $\forall x \forall y ((x = y \wedge \varphi(x)) \rightarrow \varphi(y))$
2. $\exists x \varphi(x) \wedge \forall y \forall z ((\varphi(y) \wedge \varphi(z)) \rightarrow y = z) \rightarrow \exists x (\varphi(x) \wedge \forall y (\varphi(y) \rightarrow y = x))$

ntd.13 Soundness with Identity predicate

fol:ntd:sid:
sec

Proposition ntd.32. *Natural deduction with rules for $=$ is sound.*

Proof. Any **formula** of the form $t = t$ is valid, since for every **structure** \mathfrak{M} , $\mathfrak{M} \models t = t$. (Note that we assume the term t to be closed, i.e., it contains no variables, so variable assignments are irrelevant).

Suppose the last inference in a **derivation** is $=$ Elim, i.e., the **derivation** has the following form:

$$\frac{\begin{array}{c} \Gamma_1 \\ \vdots \\ \delta_1 \\ \vdots \\ t_1 = t_2 \end{array} \quad \begin{array}{c} \Gamma_2 \\ \vdots \\ \delta_2 \\ \vdots \\ \varphi(t_1) \end{array}}{\varphi(t_2)} =\text{Elim}$$

The premises $t_1 = t_2$ and $\varphi(t_1)$ are **derived** from **undischarged** assumptions Γ_1 and Γ_2 , respectively. We want to show that $\varphi(t_2)$ follows from $\Gamma_1 \cup \Gamma_2$. Consider a **structure** \mathfrak{M} with $\mathfrak{M} \models \Gamma_1 \cup \Gamma_2$. By induction hypothesis, $\mathfrak{M} \models \varphi(t_1)$ and $\mathfrak{M} \models t_1 = t_2$. Therefore, $\text{Val}^{\mathfrak{M}}(t_1) = \text{Val}^{\mathfrak{M}}(t_2)$. Let s be any variable assignment, and $m = \text{Val}^{\mathfrak{M}}(t_1) = \text{Val}^{\mathfrak{M}}(t_2)$. By ??, $\mathfrak{M}, s \models \varphi(t_1)$ iff $\mathfrak{M}, s[m/x] \models \varphi(x)$ iff $\mathfrak{M}, s \models \varphi(t_2)$. Since $\mathfrak{M} \models \varphi(t_1)$, we have $\mathfrak{M} \models \varphi(t_2)$. \square

Photo Credits

Bibliography